

Physical Therapy Utilization of Graded Exposure for Patients With Low Back Pain

Psychological factors are commonly used to explain the development and maintenance of chronic low back pain (LBP).^{38,39} In fact, individual studies suggest that psychological factors are stronger than physical and demographic factors for predicting LBP outcomes.^{8,20} This clinical commentary focuses on the fear-avoidance model of musculoskeletal pain (FAM), which was reviewed by Leeuw et al in 2007.²⁹ Briefly, the FAM proposes that the primary affective and cognitive components influencing pain perception are anxiety, pain-related fear (including fear of movement and reinjury), and pain catastrophizing.⁴⁸ These factors interact to determine the individual's initial behavioral response to acute pain, which occurs on a continuum from avoidance to confrontation. A confrontation behavioral response (associated with low levels of anxiety, fear, and catastrophizing) is hypothesized to be an adaptive response to pain, as the likely consequences

are return to normal vocation and social activities.³² An avoidance behavioral response (associated with high levels of anxiety, fear, and catastrophizing) is consistent with increased attention to pain and is hypothesized to be a maladaptive response.³² Long-term avoidance behavior has been hypothesized to have psychological (hyperalgesia and depression),

physical (limitations in physical performance and disuse syndrome), and societal (chronic disability) consequences.^{6,27,32,48}

The FAM was selected as the focus of this manuscript, because the model appears to have clinical relevance for patients with LBP. Specifically, previous studies suggest that questionnaires based on the FAM accurately identify poor prognosis for patients with LBP.^{1,18} Furthermore, effective LBP intervention strategies based on the FAM have been reported.^{9,22,34,46} Theoretically, the common goal of FAM-based interventions is to encourage a confrontation response in patients with LBP who normally would utilize an avoidance response. This clinical commentary focuses on the application of graded exposure, which is an FAM-based intervention that potentially has application for physical therapists treating patients with LBP.

Graded exposure intervention encourages a confrontation response by exposing patients to specific situations of which they are fearful during rehabilitation.^{46,47} Philips³⁷ originally suggested graded exposure as a treatment option for LBP, because it provided the patient with direct experience that performing fearful activities will not necessarily cause further harm by damaging the spine. Patient education is another important part of graded exposure, as it serves as a way to

• **SYNOPSIS:** The fear-avoidance model of musculoskeletal pain suggests that elevated pain-related fear is a precursor to chronic low back pain. Recent prospective studies support the predictive validity of this model, and treatment approaches based on the model have also been reported in the literature. Graded exercise/activity is one treatment approach that has been well described in the literature, with reports describing physical-therapy-specific application. Graded exposure is another intervention with the potential to reduce pain-related fear, yet physical-therapy-specific application of graded exposure has not been widely described in the literature. The

purpose of this clinical commentary was to provide information on the theoretical aspects of graded exposure, to briefly review available evidence for graded exposure, and to describe physical therapy application of graded exposure for 2 patients enrolled in a physical therapy clinical trial.

• **LEVEL OF EVIDENCE:** Therapy, level 5. *J Orthop Sport Phys Ther* 2009;39(7):496-505. doi:10.2519/jospt.2009.2983

• **KEY WORDS:** behavioral intervention, biopsychosocial, disability, fear-avoidance, kinesophobia, secondary prevention

¹Assistant Professor, Brooks Center for Rehabilitation Studies, Department of Physical Therapy, University of Florida, Gainesville, FL. ²Staff Physical Therapist, Shands Rehab at the University of Florida, Gainesville, FL. Dr Steven Z. George was supported by NIH-NIAMS grant AR051128 while preparing this manuscript. Institutional Review Board approval was granted by the University of Florida. Address correspondence to Dr Steven Z. George, PO Box 100154, University of Florida, Gainesville, FL 32611-0154. E-mail: szgeorge@phhp.ufl.edu

reduce pain-related fear and threat associated with LBP, as well as to inform the patient on the potential problems associated with long-term avoidance behavior. The actual exposure itself is done in a hierarchical fashion, and the first step is to identify specific activities feared by the patient. These activities are started at a diminished level that elicits minimal amounts of fear and then gradually increased to situations that elicit larger amounts of fear. As will be outlined later in this commentary, subsequent activity progression is based on fear ratings of the selected activities. Theoretically, successful application of graded exposure results in gradual and increasing exposure to fearful activities, and this exposure provides the patient with experiential evidence that the previously held negative beliefs were exaggerated. Ideally, this positive shift in beliefs results in improved functional performance and decreases the probability of developing chronic LBP.²⁹

Graded exercise is a behavioral intervention utilized by physical therapists and, although not a focus of this commentary, it does serve as a good contrast for graded exposure. Graded exercise is consistent with the FAM because it encourages confrontation by improving patients' exercise or activity tolerance.¹⁶ However, graded exercise progression is not based on what the patient is fearful of; instead, it is based on whether the patient meets predetermined quota levels set by the clinician. A few studies suggest that patients with LBP benefited when standard rehabilitation approaches were supplemented with graded exercise,^{22,34} and a case report detailing physical therapy management of a patient with graded exercise has also been published.²¹ Therefore, there appears to be adequate information to allow for clinical implementation of graded exercise by physical therapists.

In contrast, there is less information available for physical therapy implementation of graded exposure. LBP rehabilitation based on graded exposure principles

is consistent with the FAM, and there is potential for physical therapists to utilize graded exposure while treating patients with LBP, especially when elevated levels of pain-related fear are present.²⁹ Previous reports of graded exposure have been primarily in chronic pain settings, with treatment directed by psychologists.^{46,47} To the best of our knowledge, physical-therapy-specific implementation of graded exposure has not been reported in the literature. Therefore, the purpose of this clinical commentary is to review the available evidence for graded exposure and to describe physical therapy application of graded exposure for 2 patients enrolled in a clinical trial.

Evidence Supporting Graded Exposure

Evidence for application of graded exposure can be found from 2 sources. The first source is a series of single-subject experimental designs involving within-group comparisons for patients with widespread chronic musculoskeletal pain.^{12,46,47} In these studies, consecutive patients were treated with either graded exercise or graded exposure, and, in general, superior outcomes were associated with graded exposure for pain intensity, pain-related fear, and disability.^{12,46,47} One of the interesting aspects of these studies is that when timing of graded exercise and graded exposure was varied, graded exposure was associated with larger improvements in pain-related fear.^{46,47} Furthermore, when education about exposure was compared to actual in vivo graded exposure (ie, actually participating in exposure), graded exposure was necessary to observe increased activity.¹²

The second source of evidence for graded exposure is from 3 recently reported randomized trials that involved between-group comparisons. In 2008, Linton et al³⁵ reported the results of a clinical trial that compared the effects of graded exposure plus usual care to a wait list control group. This trial included 46 patients with at least 3 months of activity restriction due to LBP and with elevated

levels of pain-related fear. Immediately after treatment, patients receiving graded exposure in addition to usual care had statistically larger improvements in function (effect size, 0.6) but not for pain intensity or pain-related fear. This study utilized a crossover design, and when the wait list control group received graded exposure, additional statistical improvements were observed for function and pain-related fear, but not for pain intensity.

In 2008, Leeuw et al³¹ reported a trial that compared the effects of graded exercise and graded exposure. This trial included 85 patients with at least 3 months of LBP and with moderate levels of pain-related fear. At the 6-month follow-up the authors reported no statistically significant differences between graded exercise and graded exposure for functional disability, main complaints, daily activity levels, or pain intensity. The authors did note some favorable trends for graded exposure, as it was associated with larger raw improvements in functional disability and main complaints, as well as statistically significant reductions in pain catastrophizing and perceived harmfulness of activities.³¹ In a planned subgroup analysis, Leeuw et al³¹ reported that the effects of graded exposure were equivalent for patients with elevated pain-related fear.

In 2008, George et al²³ reported a trial that compared the effects of treatment-based classification physical therapy to treatment-based classification physical therapy augmented with graded exercise or graded exposure. This trial included 108 patients with an LBP duration of 24 weeks or less and all levels of pain-related fear. At the 6-month follow-up the authors reported no statistically significant differences between the 3 groups for pain and disability outcomes. However, treatment-based classification physical therapy alone and augmented with graded exposure was associated with larger 6-month improvements in pain-related fear in comparison to physical therapy augmented with graded exercise. In a planned subgroup analysis, George et al²³ reported that the effects of graded expo-

sure were equivalent for patients with elevated pain-related fear.

The available data from single-subject studies^{12,46,47} and randomized clinical trials^{23,31,35} allow some tentative conclusions to be drawn regarding evidence for graded exposure. First, consistent evidence supporting use of graded exposure comes from single-subject studies involving patients with chronic musculoskeletal pain conditions.^{12,46,47} Second, there is some preliminary randomized trial evidence suggesting that for patients with chronic LBP graded exposure is superior to usual care while waiting to receive exposure,³⁵ and weaker evidence suggesting that graded exposure is superior to graded exercise.³¹ Third, there is only 1 trial investigating graded exposure for individuals with acute and subacute LBP, and it suggests no additional benefit from graded exposure in addition to treatment-based classification physical therapy.²³ Last, there is no evidence to suggest that patients with elevated pain-related fear benefit more from graded exposure in comparison to those with normal levels of pain-related fear.^{23,31}

Physical Therapy Application of Graded Exposure

The literature investigating graded exposure is small and, obviously, additional research is needed before definitive clinical recommendations can be made. However, the available evidence does suggest the potential for physical therapy application of graded exposure and, currently, physical-therapy-specific examples of graded exposure are lacking in the literature. In this clinical commentary the application of graded exposure will be done by describing treatment details for 2 patients who participated in the previously mentioned George et al²³ clinical trial. These particular patients were selected because they received their treatment from the same clinic and were both treated by the same physical therapist. Such details are not typically provided when randomized trials are reported, and the information provided in this commentary may allow

TABLE 1			EXAMINATION FINDINGS AND TREATMENT-BASED CLASSIFICATION DETERMINATION		
Examination			Patient 1		Patient 2
Duration of low back pain (d)			84		18
Onset of low back pain			Sudden		Sudden
Symptom location			Low back and buttock		Low back and buttock
Prior history of low back pain			No		No
Lumbar movement testing (deg, symptom status)					
Total flexion			20°, status quo		60°, status quo
Total extension			20°, status quo		10°, status quo
Left side-bending			10°, status quo		22°, status quo
Right side-bending			10°, status quo		20°, status quo
Qualitative assessment of lumbar flexion			Deviation present		Deviation present
Straight-leg raise (deg, symptom reproduction)					
Right			70°, negative		76°, negative
Left			85°, negative		70°, negative
Bilateral active straight-leg raise			Success		Success
Active sit-up test			Unsuccessful		Success
Spinal tenderness			Present		Absent
Prone hip internal rotation (deg)					
Right			25°		50°
Left			25°		48°
Lumbar mobility			Hypermobility		Hypermobility
Prone instability test			Positive		Positive
Treatment classification			Stabilization		Stabilization

physical therapists to implement graded exposure in their clinical environment.

All patients enrolled in the clinical trial were evaluated using a previously described treatment-based classification system for LBP,¹³ and, because this system has been well described in the literature, this clinical commentary will not provide details on that aspect of the patient encounter. Briefly, patients underwent a focused, standard examination of the musculoskeletal system that included select historical and physical factors consistent with current trends in treatment-based classification (TABLE 1).¹⁷ The examination allowed for consideration of directional preference of lumbar spine movements,³⁶ a preliminary clinical prediction rule for stabilization,²⁴ and a validated clinical prediction rule for manipulation.¹⁰ Specifically, these 2 patients were found to have clinical signs associated with stabilization classification. Therefore, they were prescribed lumbar stabilization exercises used by Hicks et

al²⁴ in addition to graded exposure.

Clinical assessment of patients in this trial also considered patient affect and cognition style consistent with the FAM.²⁹ Pain-related fear was quantified with 2 measures, and the first was the Fear-Avoidance Beliefs Questionnaire (FABQ).⁴⁹ The FABQ was developed as a disease-specific measure of patient beliefs regarding the effect of physical activity and work on LBP. The FABQ also assesses patient beliefs regarding reinjury and return to work.⁴⁹ Individual FABQ items are scored from 0 to 6, and the 4-item physical activity scale (FABQ-PA) has a total range of 0 to 24, while the 7-item work scale (FABQ-W) has a total range of 0 to 42. Separate consideration of the physical activity and work domains are recommended, with higher scores indicating higher fear-avoidance levels for both scales.⁴⁹ The FABQ has demonstrated adequate reliability and validity in previous studies.^{28,43–45,49}

To implement graded exposure treat-

TABLE 2

FEAR OF DAILY ACTIVITIES RATING
RESPONSES FOR PATIENT 1

Measure	Initial Rating	4-wk Rating
FABQ physical activity scale score (0-24)	21	13
Fear of Daily Activities Questionnaire (FDAQ)		
Sitting for longer than 1 h*	20	10
Standing for longer than 30 min*	5	0
Walking for longer than 30 min*	25	10
Lifting less than 9.1 kg*	10	10
Lifting more than 9.1 kg*	40	30
Carrying less than 9.1 kg*	10	10
Carrying more than 9.1 kg*	40	30
Twisting* [†]	40	10
Reaching to floor* [†]	50	15
Performing back exercises*	10	5

Abbreviation: FABQ, Fear-Avoidance Beliefs Questionnaire.

* Close-ended item (range, 0-100) from FDAQ.

[†] Activity was used for graded exposure treatment.

TABLE 3

FEAR OF DAILY ACTIVITIES RATING
RESPONSES FOR PATIENT 2

Activity	Initial Rating	4-wk Rating
FABQ physical activity scale score (0-24)	19	1
Fear of Daily Activities Questionnaire (FDAQ)		
Sitting for longer than 1 h*	50	0
Standing for longer than 30 min*	50	0
Walking for longer than 30 min*	60	0
Lifting less than 9.1 kg*	20	0
Lifting more than 9.1 kg*	40	5
Carrying less than 9.1 kg*	20	0
Carrying more than 9.1 kg*	40	5
Twisting*	10	0
Reaching to floor*	20	0
Performing back exercises*	30	0
Walking up steep hill ^{†‡}	60	0
Folding laundry while standing ^{†‡}	60	0

Abbreviation: FABQ, Fear-Avoidance Beliefs Questionnaire.

* Close-ended item from FDAQ.

[†] Open-ended item from FDAQ.

[‡] Activity was used for graded exposure treatment.

therapist then identifies items ranked as equal to or higher than 40/100 for implementation by graded exposure. This process is a modification of a previously reported method using photographs to select highly rated fear activities for inclusion in graded exposure treatment.³⁰ This modification was necessary because use of photographs was considered too time consuming for the clinical environment in which most physical therapists practice. It is important to note that while the FDAQ is utilized in our clinical setting for purposes of determining fear of activities, its psychometric properties have not been reported in the peer-review literature.

The 2 patients highlighted in this clinical commentary completed the FABQ and the FDAQ (TABLES 2 and 3). The FABQ scores were used to get a general indication of the levels of pain-related fear for these patients. In this particular clinical setting, the FABQ-PA was more appropriate for scoring because these patients did not have work-related injury. Although there is not a widely accepted threshold for determining elevated FABQ-PA scores, each of these patients exceeded the median score reported by Burton et al.⁹ The FDAQ results were considered for decision making related to graded exposure intervention.

Patient 1 responded to the 10 standard FDAQ items and did not provide any open-ended responses. The FDAQ showed variability in fear ratings of specific activities. The items corresponding to twisting (40/100) and reaching to the floor (50/100) had the highest fear ratings and were considered for graded exposure treatment. Patient 2 responded to the 10 standard FDAQ items, and provided 2 open-ended responses: fear of walking up a steep hill and folding laundry while standing. Patient 2 also had variability in fear ratings and the open-ended responses corresponded with the highest pain ratings (60/100). These activities were considered for the graded exposure treatment. It was hypothesized that a decrease in self-report of disability and pain-related fear would be associated

ment, fear related to specific activities must be assessed, in addition to general pain-related fear levels. Fear of specific activities was determined using an idiosyncratic clinical questionnaire, called the Fear of Daily Activities Questionnaire (FDAQ). The FDAQ lists 10 activities that patients with LBP commonly report being fearful of performing due

to pain (APPENDIX). The FDAQ also has 2 options for open-ended responses, so that patients with LBP can provide additional examples and ratings of activities that they fear performing due to pain. Patients rate each FDAQ item using a numerical rating scale that ranges from 0 (no fear) to 100 (maximal fear).

After the items are rated, the physical

[CLINICAL COMMENTARY]

with successful graded exposure treatment for these patients.

INTERVENTION

GRADED EXPOSURE WAS ADMINISTERED following suggestions provided from available studies^{12,46,47} and reviews on the FAM.^{29,48} One component of the treatment is to provide patient education that decreases fear and threat associated with LBP. Another component of the treatment is to implement activities that were rated as fearful under physical therapist supervision. In clinical situations, these components are implemented simultaneously, but for clarity they will be described separately in this clinical commentary.


Pain-related fear associated with LBP can be positively influenced by the type of information patients receive from their healthcare provider. The *Back Book*⁴² is one way this information can be delivered. The *Back Book* provides current theory and evidence associated with the management of LBP in a factual, straightforward, and nonthreatening manner. Key principles of the *Back Book* are summarized in **TABLE 4** and randomized clinical trials using the *Back Book* have demonstrated that it effectively reduced FABQ scores, either alone^{7,9} or in combination with physical therapy.²² There is also evidence to suggest that use of the *Back Book* is especially warranted for patients with elevated levels of pain-related fear.^{9,22}

Both patients described in this commentary were issued and asked to read the *Back Book* pamphlet. The physical therapist then assessed whether the pamphlet was read, and encouraged discussion of key principles from the *Back Book*. For example, patient 2 had questions about why LBP is not always an indication of damage to the spine. This provided an opportunity for the physical therapist to review with the patient how radiological indicators of damage to the spine (ie, herniated nucleus pulposus, intervertebral disc

degeneration, spondylolysis, and spondylolisthesis) are common in patients with and without LBP.³⁻⁵

Progression or modification of this educational message does not typically occur during subsequent treatment ses-

TABLE 4		KEY PRINCIPLES FROM <i>BACK BOOK</i> EDUCATIONAL PAMPHLET	
Category		Principle	
Abnormal imaging findings		• Very rarely a sign of serious disease • Commonly found in people without low back pain	
Implications of low back pain		• No suggestion of permanent damage • The spine is strong, even when it is painful • Pain does not mean your back has serious damage	
Treatment of low back pain		• A number of treatments can help to control the pain • Lasting relief depends on your effort • Concentrate on maintaining and improving activity to restore normal function and fitness • Utilize positive attitude and adaptive coping skills	



sions. Instead the physical therapist takes the opportunity to reinforce key principles from the *Back Book*. For example, patient 1 had questions about why performing fearful activities would assist with his recovery. This provided an opportunity for the physical therapist to review the FAM with the patient and to indicate how continued avoidance of fearful activities could potentially result in the development of chronic pain and disability.

Graded exposure to activities rated as highly fearful were included in the treatment plan, using parameters described in **FIGURE 1**.²³ During the first session, 2 highly rated activities from the FDAQ were introduced to the patient at intensities, frequencies, and durations that did not increase fear. This initial level of activity was determined through patient history or by having the patient perform the activity at a self-selected level that did not increase fear. Subsequent patient activity was progressed if decreased fear was reported following exposure to the fearful activity (**FIGURE 1**). When decreased fear of an activity was reported, the patient received positive reinforcement for confrontation. Then the intensity, frequency, and/or duration of activity increased for the subsequent session. When maintained or increased fear of activity was reported, the importance of attempting these activities was explained to the patient. Then the activity exposure was maintained at the current intensity, frequency, and duration for the subsequent session. The decision to increase or maintain exposure levels is made from ratings that occur within the same treatment session (**FIGURE 1**), not in ratings that occur between treatment sessions.²³ Once fear levels are decreased, patients are also encouraged to expose themselves to these activities outside of the clinic, as part of the home exercise program; although it should be noted that compliance to the home exercise program portion of graded exposure was not measured for these patients.²³

Because it is not feasible to describe each treatment session for both patients,

TABLE 5		TREATMENT SUMMARY FOR INITIAL GRADED EXPOSURE TREATMENT SESSIONS FOR PATIENT 2		
	Physical Therapy Session 1	Physical Therapy Session 2	Physical Therapy Session 3	
Fearful activity 1: walking up steep hill				
Preactivity fear level	60/100	20/100	20/100	
Clinic activity	Treadmill walking, 3 min, 3.0% incline, 1.9 km/h	Treadmill walking, 4 min, 3.0% incline, 1.9 km/h	Treadmill walking, 4 min, 3.0% incline, 1.9 km/h	
Postactivity fear level	10/100	30/100	10/100	
Fear reduced?	Yes	No	Yes	
Plan	Progress exposure time	Maintain exposure level	Progress exposure time and incline	
Fearful activity 2: folding laundry				
Preactivity fear level	60/100	50/100	30/100	
Clinic activity	Floor-to-table lift, 4.5 kg, 10 times	Floor-to-table lift, 4.5 kg, 12 times; standing assembly activity, 5 min	Floor-to-table lift, 6.8 kg, 10 times; standing assembly activity, 7 min	
Postactivity fear level	20/100	30/100	20/100	
Fear reduced?	Yes	Yes	Yes	
Plan	Increase repetition and include standing time	Increase lifting weight and standing time	Increase repetition and standing time	

only the first 3 graded exposure treatment sessions for patient 2 are summarized (**TABLE 5**). The decision-making process for the first 3 sessions of patient 2 provide an adequate demonstration of the application of graded exposure treatment. These examples will allow readers to become familiar with typical decision-making patterns involved with graded exposure treatment.²³

Patient 2 used the open-ended questions of the FDAQ to indicate highly rated fearful activities for walking up a steep hill and folding laundry while standing (**TABLE 3**). The fear of walking up a steep hill was initially rated by the patient at 60/100. The physical therapist used information from the patient history (walking time limit) and from having the patient walk on the treadmill in the clinic (incline level and walking speed) to determine initial activity level during session 1 (**TABLE 5**). The patient performed the treadmill walking as suggested and reported a postactivity fear rating of 10/100. At the conclusion of session 1, the physical therapist planned to increase

exposure time for the next treatment session. During session 2, the patient rated preactivity fear of walking up a steep hill at 20/100 (**TABLE 5**). Although this was an increase from the postactivity rating from session 1 (10/100), the therapist still increased the exposure time because the patient had previously noted a decrease during session 1. The patient completed 4 minutes on the treadmill as suggested and reported a postactivity fear rating of 30/100. At the conclusion of session 2, the physical therapist planned to maintain the exposure level due to this increase in fear rating. Session 3 began with a preactivity fear rating of 20/100. Again, even though this represented a decrease in fear from the postactivity rating from session 2 (30/100), the previous exposure level was maintained, because the patient had reported an increase in fear following the activity during session 2. The patient completed the suggested walking activity in session 3 and rated the postexercise fear rating at 10/100. At the conclusion of session 3, the physical therapist made a plan to increase the

walking time and incline for session 4. The graded exposure progression for inclined walking continued in a similar fashion until the patient was walking 10 minutes at 2.9 km/h on a 3.5% incline, with preactivity and postactivity fear ratings of 0/100.

Folding laundry while standing was the second activity rated as highly fearful (60/100). After further questioning from the physical therapist, it was determined that there were 2 parts of this activity that caused the patient fear. The first part was lifting the basket when it was full of clothes and the second part was standing in place to fold the laundry. Therefore, the physical therapist implemented both activities during the graded exposure treatment component. The physical therapist used information from the patient history (height of lift) and from having the patient practice lifts (weight of lift) to determine initial activity level during session 1 (TABLE 5). The preactivity to postactivity fear ratings decreased from 60/100 to 20/100 following session 1. At the conclusion of session 1, the physical therapist planned to increase exposure by increasing the number of repetitions and by adding a standing activity to the treatment program. During session 2, the patient rated fear from folding laundry at 50/100 (TABLE 5). Anecdotally, the patient also reported some anxiety about performing the standing activity after the increase in lifting repetitions. Although this was an increase from the postactivity fear rating from session 1 (20/100), the therapist still increased the exposure as planned from the decrease during session 1. The physical therapist acknowledged the patient's fear and anxiety, but also let the patient know that her fear and anxiety were the reasons for selecting these particular activities. The patient was able to perform the activities as prescribed during session 2, and fear ratings were decreased from 50/100 to 30/100. At the conclusion of session 2, the physical therapist planned to increase exposure by increasing the weight of the lift and by increasing the

TABLE 6		SUMMARY OF OUTCOMES FOLLOWING GRADED EXPOSURE	
Measure	Initial Rating	4-wk Rating	
Patient 1			
Disability, ODQ (0%-100%)	50%	14%*	
Average pain intensity (at present), ordinal scale (0-10)	7/10	6/10	
Pain (at worst), ordinal scale (0-10)	7/10	6/10	
Pain (at best), ordinal scale (0-10)	5/10	3/10*	
Patient 2			
Disability, ODQ (0%-100%)	46%	20%*	
Pain intensity (at present), ordinal scale (0-10)	4/10	2/10*	
Pain (at worst), ordinal scale (0-10)	8/10	3/10*	
Pain (at best), ordinal scale (0-10)	2/10	2/10	
Abbreviation: ODQ, Oswestry Disability Questionnaire.			
* Improvement met criterion for minimally clinically important difference (6 points or greater on ODQ and 2 points or greater for pain ratings).			

time of the standing activity. Session 3 began with a pre-exercise activity fear rating of 30/100, and the patient was able to perform all the exposure-related activity, with postactivity fear ratings of 10/100. This graded exposure progression for folding laundry while standing continued in a similar fashion until the patient was lifting 9.9 kg 15 times and doing standing activities for 15 minutes, with preactivity and postactivity fear ratings of 0/100.

OUTCOMES

STANDARDIZED OUTCOME MEASURES were used to assess patient status before treatment, and then 4 weeks later.²³ Self-reported disability was assessed with the Modified Oswestry Disability Questionnaire (ODQ).¹⁵ The ODQ has been found to have high levels of test-retest reliability, validity, and responsiveness,^{15,19,41} and is recommended as a primary outcome measure of self-report of disability for patients with LBP.¹⁴ Pain intensity was assessed using an numerical rating scale (range, 0-10), and the patient was asked to rate the current, worst (highest), and best (lowest) pain intensity ratings over the past 24 hours. The mean of these ratings was reported in the clinical trial.²³ This method of pain assessment has

previously demonstrated acceptable reliability and validity in the literature.^{25,40} Although not commonly utilized as outcome measures, the FABQ and the FDAQ were also collected 4 weeks later.

Patient 1 had 4 treatments that occurred over the 4-week period, and changes in status are summarized in TABLES 2 and 6. The observed improvement in pain-related disability exceeded a published threshold for minimally clinically important difference (MCID).¹⁹ The observed improvement in pain intensity met the MCID threshold for one third of the ratings.¹¹ Patient 1 also reported improvements in pain-related fear, and fear of activities included in the graded exposure program (twisting and reaching). Patient 2 had 9 treatments over the 4-week period, with improvements that exceeded the MCID for the modified ODQ,¹⁹ and met the MCID threshold for two thirds of the pain intensity ratings (TABLES 2 and 5).¹¹ Patient 2 reported improvements in pain-related fear, and fear of activities included in the graded exposure program (walking up a steep hill and folding laundry while standing). Patient 2 had larger improvements in pain-related fear and fear of specific activities in comparison to patient 1. However, it is important to note that there are no accepted MCIDs for the FABQ and FDAQ, so determining

the clinical relevance of these improvements is difficult.

An important issue is whether the 2 patients highlighted in this clinical commentary were representative of the graded exposure outcomes from the overall randomized trial. To address this issue, the 4-week disability and pain intensity outcomes for all patients receiving graded exposure were placed in a graph (FIGURE 2).²³ The average 4-week change in the modified ODQ scale was 13.0 (SD, 14.8), so outcomes for patient 1 were within 2 standard deviations of the mean, and outcomes for patient 2 were within 1 standard deviation of the mean (FIGURE 2A). The average 4-week change in average pain intensity was 2.7 (SD, 2.2), so outcomes for patients 1 and 2 were within 1 standard deviation of the mean (FIGURE 2B). Therefore, the patients highlighted in this commentary were representative of overall trial outcomes. However, it should also be recognized that expected outcomes are likely to change based on initial levels of disability and fear. Furthermore, change in ratings for single patients must be viewed with caution, as they do not control for measurement variation and nonspecific effects. As with any treatment approach, variability in response to graded exposure is expected, and only the randomized trial can provide a more definitive assessment of the effectiveness of the described approach.²³ The patients in this commentary were neither the best nor worst responders and were used only to provide an example of a standard approach to graded exposure, not to validate the effectiveness of graded exposure.

SUMMARY

GRADED EXPOSURE IS A THEORETICALLY appropriate way to reduce chronic pain and disability within the FAM.²⁹ The primary goal of graded exposure is behavioral in nature, as it serves as a way to increase the performance of fearful activities. There are also important cognitive aspects of applying




FIGURE 2. (A) Change in disability scores for patients receiving graded exposure intervention in physical therapy clinical trial at 4 weeks.²⁵ (B) Change in average pain intensity ratings for patients receiving graded exposure intervention in physical therapy clinical trial at 4 weeks.²⁵

graded exposure that shouldn't be overlooked. Patients receive education that decreases the fear and threat associated with LBP. Also, patients receive positive reinforcement for performing fearful activities and utilizing beneficial coping strategies. Therefore, it is impossible to separate the effects of the education and activity implementation components of the graded exposure treatment. In our opinion it is best to view graded exposure as a combined cognitive and behavioral treatment that encourages confrontation of fearful activities during rehabilitation of LBP.

Utilizing graded exposure may represent a philosophical shift in how physical therapists typically manage LBP.

For example, the goal of patient education changes when using behavioral approaches. Instead of providing purely mechanical or anatomical explanations for causes and treatment of LBP, anatomical explanations of LBP are downplayed and use of adaptive coping strategies is encouraged. Furthermore, exercise prescription for LBP is typically based on physiological (muscle force production, range of motion, flexibility, or aerobic capacity), temporal (chronic or acute symptoms), or anatomical location (leg pain or no leg pain) parameters.^{2,33} In contrast, exercise prescription for graded exposure is based on identifying fearful activities, systematically exposing the patients to these fearful activities, and progressing exposure to the activities when fear ratings decreased.^{23,29,31,35}

Physical therapy practice guidelines allow for the assessment of psychological factors that influence rehabilitation outcomes, although such assessment is not commonly performed by clinicians.^{2,26,33} The FAM is a specific theoretical model that provides guidance for clinicians to identify patients with elevated pain-related fear and appropriately modify treatment. This clinical commentary described physical therapy utilization of graded exposure, which has not been widely reported in the literature. Given there is limited evidence currently available on graded exposure, additional research from prognostic studies and clinical trials is necessary to guide future clinical application of graded exposure for physical therapists. High-priority items for future research in graded exposure include confirming whether graded exposure is superior to other behavioral interventions for chronic LBP and whether subgroups of patients that will receive additional benefit from graded exposure can be accurately identified to allow focused application of the intervention. ●

ACKNOWLEDGEMENT: We thank Shands Rehab at the University of Florida for utilization of facilities.

REFERENCES

1. Al-Obaidi SM, Beattie P, Al-Zoabi B, Al-Wekeel S. The relationship of anticipated pain and fear avoidance beliefs to outcome in patients with chronic low back pain who are not receiving workers' compensation. *Spine*. 2005;30:1051-1057.
2. Battie MC, Cherkin DC, Dunn R, Ciol MA, Wheeler KJ. Managing low back pain: attitudes and treatment preferences of physical therapists. *Phys Ther*. 1994;74:219-226.
3. Beutler WJ, Fredrickson BE, Murtland A, Sweeney CA, Grant WD, Baker D. The natural history of spondylolysis and spondylolisthesis: 45-year follow-up evaluation. *Spine*. 2003;28:1027-1035; discussion 1035. <http://dx.doi.org/10.1097/01.BRS.0000061992.98108.A0>
4. Boden SD, Davis DO, Dina TS, Patronas NJ, Wiesel SW. Abnormal magnetic-resonance scans of the lumbar spine in asymptomatic subjects. A prospective investigation. *J Bone Joint Surg Am*. 1990;72:403-408.
5. Boos N, Semmer N, Elfering A, et al. Natural history of individuals with asymptomatic disc abnormalities in magnetic resonance imaging: predictors of low back pain-related medical consultation and work incapacity. *Spine*. 2000;25:1484-1492.
6. Bortz WM, 2nd. The disuse syndrome. *West J Med*. 1984;141:691-694.
7. Buchbinder R, Jolley D, Wyatt M. 2001 Volvo Award Winner in Clinical Studies: Effects of a media campaign on back pain beliefs and its potential influence on management of low back pain in general practice. *Spine*. 2001;26:2535-2542.
8. Burton AK, Tilotson KM, Main CJ, Hollis S. Psychosocial predictors of outcome in acute and subchronic low back trouble. *Spine*. 1995;20:722-728.
9. Burton AK, Waddell G, Tilotson KM, Summerton N. Information and advice to patients with back pain can have a positive effect. A randomized controlled trial of a novel educational booklet in primary care. *Spine*. 1999;24:2484-2491.
10. Childs JD, Fritz JM, Flynn TW, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med*. 2004;141:920-928.
11. Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine*. 2005;30:1331-1334.
12. de Jong JR, Vlaeyen JW, Onghena P, Goossens ME, Geilen M, Mulder H. Fear of movement/(re) injury in chronic low back pain: education or exposure in vivo as mediator to fear reduction? *Clin J Pain*. 2005;21:9-17; discussion 69-72.
13. Delitto A, Erhard RE, Bowling RW. A treatment-based classification approach to low back pain syndrome: identifying and staging patients for conservative treatment. *Phys Ther*. 1995;75:470-485; discussion 485-479.
14. Deyo RA, Battie M, Beurskens AJ, et al. Outcome measures for low back pain research. A proposal for standardized use. *Spine*. 1998;23:2003-2013.
15. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine*. 2000;25:2940-2952; discussion 2952.
16. Fordyce WE, Fowler RS, Jr., Lehmann JF, Delateur BJ, Sand PL, Trieschmann RB. Operant conditioning in the treatment of chronic pain. *Arch Phys Med Rehabil*. 1973;54:399-408.
17. Fritz JM, Cleland JA, Childs JD. Subgrouping patients with low back pain: evolution of a classification approach to physical therapy. *J Orthop Sports Phys Ther*. 2007;37:290-302. <http://dx.doi.org/10.2519/jospt.2007.2498>
18. Fritz JM, George SZ. Identifying psychosocial variables in patients with acute work-related low back pain: the importance of fear-avoidance beliefs. *Phys Ther*. 2002;82:973-983.
19. Fritz JM, Irrgang JJ. A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. *Phys Ther*. 2001;81:776-788.
20. Gatchel RJ, Polatin PB, Mayer TG. The dominant role of psychosocial risk factors in the development of chronic low back pain disability. *Spine*. 1995;20:2702-2709.
21. George SZ, Bialosky JE, Fritz JM. Physical therapist management of a patient with acute low back pain and elevated fear-avoidance beliefs. *Phys Ther*. 2004;84:538-549.
22. George SZ, Fritz JM, Bialosky JE, Donald DA. The effect of a fear-avoidance-based physical therapy intervention for patients with acute low back pain: results of a randomized clinical trial. *Spine*. 2003;28:2551-2560.
23. George SZ, Zeppieri G, Jr., Cere AL, et al. A randomized trial of behavioral physical therapy interventions for acute and sub-acute low back pain (NCT00373867). *Pain*. 2008;140:145-157. <http://dx.doi.org/10.1016/j.pain.2008.07.029>
24. Hicks GE, Fritz JM, Delitto A, McGill SM. Preliminary development of a clinical prediction rule for determining which patients with low back pain will respond to a stabilization exercise program. *Arch Phys Med Rehabil*. 2005;86:1753-1762.
25. Jensen MP, Turner JA, Romano JM. What is the maximum number of levels needed in pain intensity measurement? *Pain*. 1994;58:387-392.
26. Jette AM, Smith K, Haley SM, Davis KD. Physical therapy episodes of care for patients with low back pain. *Phys Ther*. 1994;74:101-110; discussion 110-105.
27. Kottke FJ. The effects of limitation of activity upon the human body. *JAMA*. 1966;196:825-830.
28. Kovacs FM, Muriel A, Medina JM, Abaira V, Sanchez MD, Jauregui JO. Psychometric characteristics of the Spanish version of the FAB questionnaire. *Spine*. 2006;31:104-110.
29. Leeuw M, Goossens ME, Linton SJ, Crombez G, Boersma K, Vlaeyen JW. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med*. 2007;30:77-94.
30. Leeuw M, Goossens ME, van Breukelen GJ, Boersma K, Vlaeyen JW. Measuring perceived harmfulness of physical activities in patients with chronic low back pain: the Photograph Series of Daily Activities--short electronic version. *J Pain*. 2007;8:840-849. <http://dx.doi.org/10.1016/j.jpain.2007.05.013>
31. Leeuw M, Goossens ME, van Breukelen GJ, et al. Exposure in vivo versus operant graded activity in chronic low back pain patients: results of a randomized controlled trial. *Pain*. 2008;138:192-207. <http://dx.doi.org/10.1016/j.pain.2007.12.009>
32. Lethem J, Slade PD, Troup JD, Bentley G. Outline of a Fear-Avoidance Model of exaggerated pain perception--I. *Behav Res Ther*. 1983;21:401-408.
33. Li LC, Bombardier C. Physical therapy management of low back pain: an exploratory survey of therapist approaches. *Phys Ther*. 2001;81:1018-1028.
34. Lindstrom I, Ohlund C, Eek C, et al. The effect of graded activity on patients with subacute low back pain: a randomized prospective clinical study with an operant-conditioning behavioral approach. *Phys Ther*. 1992;72:279-290; discussion 291-273.
35. Linton SJ, Boersma K, Jansson M, Overmeer T, Lindblom K, Vlaeyen JW. A randomized controlled trial of exposure in vivo for patients with spinal pain reporting fear of work-related activities. *Eur J Pain*. 2008;12:722-730. <http://dx.doi.org/10.1016/j.ejpain.2007.11.001>
36. Long AL. The centralization phenomenon. Its usefulness as a predictor or outcome in conservative treatment of chronic low back pain (a pilot study). *Spine*. 1995;20:2513-2520; discussion 2521.
37. Philips HC. Avoidance behaviour and its role in sustaining chronic pain. *Behav Res Ther*. 1987;25:273-279.
38. Pincus T, Burton AK, Vogel S, Field AP. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine*. 2002;27:E109-120.
39. Pincus T, Vlaeyen JW, Kendall NA, Von Korf MR, Kalauokalani DA, Reis S. Cognitive-behavioral therapy and psychosocial factors in low back pain: directions for the future. *Spine*. 2002;27:E133-138.
40. Roach KE, Brown MD, Dunigan KM, Kusek CL, Walas M. Test-retest reliability of patient reports of low back pain. *J Orthop Sports Phys Ther*. 1997;26:253-259.
41. Roland M, Fairbank J. The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire. *Spine*. 2000;25:3115-3124.
42. Roland M, Waddell G, Klaber-Moffett J. *The Back Book*. Norwich, UK: The Stationery Office; 1996.
43. Staerkle R, Mannion AF, Elfering A, et al. Longitudinal validation of the fear-avoidance beliefs questionnaire (FABQ) in a Swiss-German

sample of low back pain patients. *Eur Spine J*. 2004;13:332-340.

44. Swinkels-Meewis IE, Roelofs J, Verbeek AL, Oostendorp RA, Vlaeyen JW. Fear of movement/(re)injury, disability and participation in acute low back pain. *Pain*. 2003;105:371-379.
45. Swinkels-Meewis IE, Roelofs J, Verbeek AL, Oostendorp RA, Vlaeyen JW. Fear-avoidance beliefs, disability, and participation in workers and non-workers with acute low back pain. *Clin J Pain*. 2006;22:45-54.
46. Vlaeyen JW, de Jong J, Geilen M, Heuts PH,

van Breukelen G. Graded exposure in vivo in the treatment of pain-related fear: a replicated single-case experimental design in four patients with chronic low back pain. *Behav Res Ther*. 2001;39:151-166.

47. Vlaeyen JW, de Jong J, Geilen M, Heuts PH, van Breukelen G. The treatment of fear of movement/(re)injury in chronic low back pain: further evidence on the effectiveness of exposure in vivo. *Clin J Pain*. 2002;18:251-261.
48. Vlaeyen JW, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain:

a state of the art. *Pain*. 2000;85:317-332.

49. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain*. 1993;52:157-168.



MORE INFORMATION
WWW.JOSPT.ORG

APPENDIX

FEAR OF DAILY ACTIVITIES QUESTIONNAIRE (FDAQ)

People with low back pain have told us that they are fearful of performing certain activities because they believe these activities will cause additional low back pain, or reinjure their back. Examples of such activities are listed below. Using the provided scale, please rate each activity for the amount of fear it causes you, as it relates to your low back pain. Because not all activities are fearful for all people, we are also asking you to list 2 different activities that cause you fear, and to rate the fear associated with those activities.

0 ————— 100
No fear of activity Maximal fear of activity

Activity	Rating (0-100)
1. Sitting for longer than 1 hour	_____
2. Standing for longer than 30 minutes	_____
3. Walking for longer than 30 minutes	_____
4. Lifting less than 20 pounds*	_____
5. Lifting 20 pounds* or more	_____
6. Carrying less than 20 pounds*	_____
7. Carrying 20 pounds* or more	_____
8. Twisting	_____
9. Reaching to the floor	_____
10. Performing back exercises	_____
11. _____	_____
12. _____	_____

The FDAQ is a preliminary instrument with undocumented reliability and validity; therefore, it should be used with caution in clinical settings. This instrument was included because it was an essential part of the decision-making process for implementation of graded exposure. Future studies will report psychometric information for the FDAQ, and this information will help to determine if the instrument is appropriate for widespread use. *Twenty pounds is approximately 9.1 kg.

NOTIFY *JOSPT* of Changes in Address

Please remember to let *JOSPT* know about **changes in your mailing address**. The US Postal Service typically will not forward second-class periodical mail. Journals are destroyed, and the USPS charges *JOSPT* for sending them to the wrong address. You may change your address online at www.jospt.org. Visit **"INFORMATION FOR READERS"**, click **"Change of Address"**, and select and complete the online form. We appreciate your assistance in keeping *JOSPT*'s mailing list up to date.